

Ichthyosis Follicularis with Alopecia and Photophobia Syndrome (IFAP): A Case Report and Review of the Literature

Case Report

Elharrouni Alaoui Aicha*, Baybay Hanane, Douhi Zakia, Elloudi Sara, and Mernissi FZ

Department of Dermatology, CHU Hassan II, Morocco

Received: May 23, 2020; **Accepted:** June 12, 2020; **Published:** June 15, 2020

***Corresponding author:** El harrouni Alaoui .A, Department of dermatology ,CHU Hassan II, FES, Morocco

Copyright: © 2020 El harrouni Alaoui. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

IFAP syndrome is a rare autosomal recessive X-linked disease characterized by the triad of alopecia universalis, severe photophobia, and follicular ichthyosis. It is caused by loss of function of the gene MBTPS2. Its severity varies and there are only a few reports in the literature. We present a patient with characteristic clinical features.

Keywords

Ichthyosis Follicularis; Atrichia; Photophobia; X-linked; MBTPS2 gene; IFAP syndrome

Introduction

IFAP syndrome is a rare genetic disorder characterized by ichthyosis and alopecia from birth and sometimes accompanied by short stature, intellectual disability, and seizures epilepsy that develop in the first few years of life. Photophobia may also be present in the first year of life or appears in infancy or early childhood. Its mode of inheritance is X-linked recessive, thus mostly affecting males. Affected or carrier females may display some of its clinical features.

Case Report

An 18-month-old boy has non consanguineous parents. Pregnancy was uneventful and the boy was born underweight and preterm at 36 weeks and 5 days by natural birth and there was no history of similar illness in the family or familiar history of atopy. He was the first birth in his family. He was referred to the dermatology department with congenital universal alopecia. The clinical examination revealed a photophobia, prominent forehead, large ears with staturponderal delay and in

the dermatological examination showed skin phototype II, spiky follicular ichthyosis on the scalp and extensor aspects of the extremities, including elbows and knees, generalized dryness of the skin, angular cheilitis, follicular papules



Figure 1: Clinical appearance and dermatologic findings of the patient. Note the alopecia with an absence of scalp hair, eyebrows, and lashes (a), (b). Spiky follicular hyperkeratosis on the trunk. Skin was dry and scaly (a), angular cheilitis (b), plantar keratoderma (c) and dystrophic nails (e),(f).

and pustules on an erythematous base scattered, plantar keratoderma, associated with universal alopecia of scalp hair, eyebrows, and eyelashes, periungueal inflammation, dystrophic nails (Figure 1).

Histopathological examination of the cutaneous biopsy showed dilated hair follicles with keratin plugs extending above the surface of the skin, decreased or absent sebaceous glands and normal sweat glands, with the presence of the rare miniaturized atrophic hair follicles including rare dystrophic hair rods, giving the appearance of follicular keratosis with non-cicatricial alopecia in the context of the IFAP syndrome. He was observed by a Medical Genetics consultant where the IFAP syndrome was retained and he underwent conservative treatment with moisturizers and topical steroids and eye lubrication, with a good improvement.

Discussion

Ichthyosis follicularis, alopecia, and photophobia (IFAP) syndrome (OMIM 308205) is an extremely rare X-linked oculocutaneous genetic disorder characterized by the triad of hyperkeratotic follicular papules, total to subtotal alopecia, and photophobia of varying degrees [1]. This association was first described as a syndrome by MacLeod in 1909 [1]. Since; only 37 cases of IFAP syndrome have been reported in the literature [2-4] and most of these had additional features including atopic eczema 36% cases [5].

In clinical presentation, Follicular ichthyosis predominantly distributed on the scalp and extensor aspects of the extremities, including elbows and knees, represents the classical cutaneous manifestations, which could give the skin "nutmeg grater-" or "sandpaper-" like texture [6]. Congenital alopecia involving the scalp, eyebrows and eyelashes is another essential cutaneous manifestation of IFAP. A non cicatricial complete body alopecia is another characteristic dermatological finding of this syndrome [6]. Variable degrees of a collodion membrane may be present in the neonate. Psoriasisiform plaques, angular cheilitis, periungueal inflammation, dystrophic nails, hypohidrosis, and atopic eczema was also described [6]. The palms and soles can be affected [7], as seen in our case. Photophobia is an essential feature for the diagnosis of IFAP. It can be present early in life or later in childhood, but other Ophthalmological manifestations have been reported as well; corneal erosions, scarring atopic keratoconjunctivitis, nystagmus, and myopia

[2] Growth retardation, represented by short stature, is often observed, as seen in our case [8,9]. Psychomotor developmental delay, seizures and hypotonia were also noted [10,11]. Nevertheless, our patient did not show neurologic involvement. Other clinical features associated with IFAP syndrome consist of short stature, dysmorphic face, intestinal renal, cardiac and vertebral abnormalities, and cleft hands have been reported [12,13]. Skin histopathology is non-specific [14,15].

IFAP syndrome results from missense mutations in the membrane-bound transcription factor protease site 2 (MBTPS2) genes. This gene encodes a zinc metalloproteinase intramembrane essential for cholesterol homeostasis and response of the endoplasmic reticulum to stress [3].

The therapeutic goal is essentially a multipronged approach that aimed to repair and protect the skin barrier, reduce microbial colonization and immune regulation to greater therapeutic efficacy, and improve quality of life of these patients. Follicular hyperkeratosis can be treated using topical keratolytics, urea preparations, and emollients. Oral retinoids at a dosage of 0.3 to 1 mg/kg/day have previously been used with moderate response. Photophobia and seizures may improve spontaneously; intensive lubrication of the ocular surface remains the mainstay of therapy, although sometimes antiseizure drugs are required [16]. For our case, the treatment was essentially based on urea emollients, local corticosteroids. Life expectancy in patients with IFAP syndrome can vary from death in the neonatal period to normal surviving. Cardiopulmonary complications remain the major cause of death [17].

Conclusion

IFAP syndrome is a rare syndrome. Its diagnosis is practically clinical. The prognosis is mainly determined by cutaneous and extracutaneous manifestations. We have reported new observation of a boy non consanguineous parents whose clinic and genetic study were the keys to diagnosis, with a good improvement under topical treatment.

References

1. MacLeod JMH. "Three cases of "ichthyosis follicularis" associated with baldness," *British Journal of Dermatology*. 1909; 21: 165-189.
2. Khandpur S, Bhat R, Ramam M. Ichthyosis follicularis, alopecia and photophobia (IFAP) syndrome treated with acitretin. *J Eur Acad Dermatol Venereol*. 2005; 19: 759-762.

3. Oeffner F, Fischer G, Happle R, König A, Betz RC, Bornholdt D, et al. IFAP syndrome is caused by deficiency in MBTPS2, an intramembrane zinc metalloprotease essential for cholesterol homeostasis and ER stress response. *Am J Hum Genet.* 2009; 84: 459–467.
4. König A, Happle R. Linear lesions reflecting lyonization in women heterozygous for IFAP syndrome (ichthyosis follicularis with atrichia and photophobia). *Am J Med Genet.* 1999; 85: 365–368.
5. M'egarbané H, M'egarbané A. "Ichthyosis follicularis, alopecia, and photophobia (IFAP) syndrome," *Orphanet Journal of Rare Diseases.* 2011; 6.
6. Chauhan S, Gahalaut P, Chandra K. Ichthyosis follicularis alopecia and photophobia syndrome: transient improvement with oral isotretinoin. *Indian J Paediatr Dermatol.* 2015; 16: 39-41.
7. Boente MC, Bibas-Bonet H, Coronel AM, Asial RA. Atrichia, ichthyosis, follicular hyperkeratosis, chronic candidiasis, keratitis, seizures, mental retardation and inguinal hernia: a severe manifestation of IFAP syndrome? *Eur J Dermatol.* 2000; 10: 98-102.
8. Cambiagli S, Barbareschi M, Tadini G. Ichthyosis follicularis with atrichia and photophobia (IFAP) syndrome in two unrelated female patients. *J Am Acad Dermatol.* 2002; 46: S156–S158.
9. Bibas-Bonet H, Fauze R, Boente MC, Asial R, et al. IFAP syndrome "plus" seizures, mental retardation, and callosal hypoplasia. *Pediatr Neurol.* 2001; 24: 228–231.
10. Eramo LR, Esterly NB, Zieserl EJ, Herrmann J. Ichthyosis follicularis with alopecia and photophobia. *Arch Dermatol.* 1985; 121: 1167–1174.
11. Martino F, D'Eufemia P, Pergola MS, Finocchiaro R, Celli M, Giampà G, et al. Child with manifestations of dermatotrichic syndrome and ichthyosis follicularis-alopecia-photophobia (IFAP) syndrome. *Am J Med Genet.* 1992; 44: 233–236.
12. M'egarbané H, Zablit C, Waked N, Lefranc G, Tomb R, M'egarbané A. Ichthyosis follicularis, alopecia, and photophobia (IFAP) syndrome: report of a new family with additional features and review. *Am J Med Genet.* 2004; 124A: 323-327.
13. Oeffner F, Fischer G, Happle R, König A, Betz RC, Bornholdt D, et al. IFAP syndrome is caused by deficiency in MBTPS2, an intramembrane zinc metalloprotease essential for cholesterol homeostasis and ER stress response. *Am J Hum Genet.* 2009; 84: 459-467.
14. M'egarbané H, M'egarbané A. Ichthyosis follicularis, alopecia, and photophobia (IFAP) syndrome, *Orphanet. J Rare Dis.* 2011; 6: 29.
15. Chauhan S, Gahalaut P, Chandra K. Ichthyosis follicularis alopecia and photophobia syndrome: transient improvement with oral isotretinoin, *Indian. J Paediatr Dermatol.* 2015; 16: 39-41.
16. Traboulsi E, Waked N, M'egarbané H, M'egarbané A. Ocular findings in ichthyosis follicularis-alopecia-photophobia (IFAP) syndrome. *Ophthalmic Genet.* 2004; 25: 153-156.
17. Keyvani K, Paulus W, Traupe H, Kiesewetter F, Cursiefen C, Huk W, et al. Ichthyosis follicularis, alopecia and photophobia (IFAP) syndrome: Clinical and neuropathological observations in a 33 year old man. *Am J Med Genet.* 1998; 78: 371-377.